

Ball Average Characterizations of Function Spaces

Dachun Yang

(Joint work)

dcyang@bnu.edu.cn

School of Mathematical Sciences

Beijing Normal University

September 17 to 23, 2017 / NPFSA-2017, Będlewo, Poland

Outline

- §I **Pointwise** characterizations of Besov and Triebel-Lizorkin spaces with smoothness not greater than 1
- §II **Ball average** characterizations of **second order** Sobolev spaces
- §III **Ball average** characterizations of **second order** Besov and Triebel-Lizorkin spaces
- §IV Further remarks

Main Motivation

- ▶ Since there exists **no differential structure** on a general metric measure space, it is still an open question how to introduce **function spaces with smoothness** on such a setting.
- ▶ Find **new characterizations** of well-known function spaces so that these new characterizations can be used as the definitions of the corresponding function spaces on **metric measure spaces**.

§I. **Pointwise** characterizations of
Besov and **Triebel-Lizorkin** spaces
with smoothness not greater than
1

Homogeneous Sobolev Spaces $\dot{W}^{m,p}(\mathbb{R}^n)$ / §I

▶ Let $m \in \mathbb{N} := \{1, 2, \dots\}$, $\mathbb{Z}_+ := \mathbb{N} \cup \{0\}$ & $p \in (1, \infty)$.

- $f \in \dot{W}^{m,p}(\mathbb{R}^n)$ (homogeneous Sobolev space) \iff
 $f \in \mathcal{S}'(\mathbb{R}^n)$ (Schwartz distribution) and $\partial^\gamma f \in L^p(\mathbb{R}^n)$ for all $|\gamma| = m$; moreover,

$$\|f\|_{\dot{W}^{m,p}(\mathbb{R}^n)} := \sum_{|\gamma|=m} \|\partial^\gamma f\|_{L^p(\mathbb{R}^n)}.$$

- **Homogeneous:** for any $\lambda \in (0, \infty)$ and $f \in \dot{W}^{m,p}(\mathbb{R}^n)$,

$$\|f(\lambda \cdot)\|_{\dot{W}^{m,p}(\mathbb{R}^n)} = \lambda^{m-n/p} \|f\|_{\dot{W}^{m,p}(\mathbb{R}^n)}.$$

Inhomogeneous Sobolev Spaces $W^{m,p}(\mathbb{R}^n)$ / §I

► Let $m \in \mathbb{N} := \{1, 2, \dots\}$, $\mathbb{Z}_+ := \mathbb{N} \cup \{0\}$ & $p \in (1, \infty)$.

- $f \in W^{m,p}(\mathbb{R}^n) \iff f \in L^p(\mathbb{R}^n)$ and $\partial^\gamma f \in L^p(\mathbb{R}^n)$ for all $|\gamma| \leq m$; moreover,

$$\|f\|_{W^{m,p}(\mathbb{R}^n)} := \sum_{|\gamma| \leq m} \|\partial^\gamma f\|_{L^p(\mathbb{R}^n)}.$$

$$\left[\|f\|_{W^{m,p}(\mathbb{R}^n)} \sim \|f\|_{L^p(\mathbb{R}^n)} + \|f\|_{\dot{W}^{m,p}(\mathbb{R}^n)} \right]$$

Besov Spaces $\dot{B}_{p,q}^\alpha(\mathbb{R}^n)$ / §I

Let $\varphi_0 \in \mathcal{S}(\mathbb{R}^n)$ (**Schwartz functions**) satisfy

$$(1.1) \quad \varphi_0(x) = 1 \text{ if } |x| \leq 1 \quad \text{and} \quad \varphi_0(y) = 0 \text{ if } |y| \geq 3/2,$$

and let

$$(1.2) \quad \varphi^{(j)}(x) := \varphi_0(2^{-j}x) - \varphi_0(2^{-j+1}x), \quad \forall x \in \mathbb{R}^n, \quad \forall j \in \mathbb{Z}.$$

Then

$$\sum_{j \in \mathbb{Z}} \varphi^{(j)}(x) = 1, \quad \forall x \in \mathbb{R}^n \setminus \{\vec{0}_n\}.$$

Let (**Triebel, 83 book**)

$$\mathcal{S}_\infty(\mathbb{R}^n) := \left\{ f \in \mathcal{S}(\mathbb{R}^n) : \int_{\mathbb{R}^n} f(x) x^\alpha dx = 0, \quad \forall \alpha \in \mathbb{Z}_+^n \right\}.$$

Besov Spaces $\dot{B}_{p,q}^\alpha(\mathbb{R}^n)$ / §I

► If $\alpha \in (0, \infty)$ & $p, q \in (0, \infty]$, then $f \in \dot{B}_{p,q}^\alpha(\mathbb{R}^n)$
(homogeneous Besov space) $\iff f \in \mathcal{S}'_\infty(\mathbb{R}^n)$ **(dual of $\mathcal{S}_\infty(\mathbb{R}^n)$)**
 such that

$$\|f\|_{\dot{B}_{p,q}^\alpha(\mathbb{R}^n)} := \left\{ \sum_{j=-\infty}^{\infty} 2^{j\alpha q} \left\| \left(\varphi^{(j)} \widehat{f} \right)^\vee \right\|_{L^p(\mathbb{R}^n)}^q \right\}^{1/q}$$

$$=: \left\| \left\{ 2^{j\alpha} \left\| \left(\varphi^{(j)} \widehat{f} \right)^\vee \right\|_{L^p(\mathbb{R}^n)} \right\}_{j \in \mathbb{Z}} \right\|_{\ell^q} < \infty.$$

► In what follows, for any function or distribution g , \widehat{g} (or g^\vee) denotes its Fourier (or **inverse Fourier**) transform.

Triebel-Lizorkin Spaces $\dot{F}_{p,q}^\alpha(\mathbb{R}^n)$ / §I

► If $\alpha \in (0, \infty)$ & $p, q \in (0, \infty]$, then $f \in \dot{F}_{p,q}^\alpha(\mathbb{R}^n)$
 (homogeneous Triebel-Lizorkin space) $\iff f \in \mathcal{S}'_\infty(\mathbb{R}^n)$ such
 that $\|f\|_{\dot{F}_{p,q}^\alpha(\mathbb{R}^n)} < \infty$, where

$$\|f\|_{\dot{F}_{p,q}^\alpha(\mathbb{R}^n)} := \left\| \left\| \{2^{j\alpha} (\varphi^{(j)} \widehat{f})^\vee\}_{j \in \mathbb{Z}} \right\|_{\ell^q} \right\|_{L^p(\mathbb{R}^n)}, \quad p < \infty$$

and

$$\|f\|_{\dot{F}_{\infty,q}^\alpha(\mathbb{R}^n)} := \sup_{\substack{x \in \mathbb{R}^n \\ m \in \mathbb{Z}}} \left\{ 2^{mn} \int_{B(x, 2^{-m})} \sum_{j=m}^{\infty} |2^{j\alpha} (\varphi^{(j)} \widehat{f})^\vee(y)|^q dy \right\}^{\frac{1}{q}}.$$

► $\dot{F}_{\infty,2}^0(\mathbb{R}^n) = \text{BMO}(\mathbb{R}^n)$, $\dot{F}_{p,2}^0(\mathbb{R}^n) = H^p(\mathbb{R}^n)$, $p \in (0, \infty)$ &
 $\dot{F}_{p,2}^m(\mathbb{R}^n) = \dot{W}^{m,p}(\mathbb{R}^n)$, $m \in \mathbb{Z}_+$, $p \in (1, \infty)$.

$B_{p,q}^\alpha(\mathbb{R}^n)$ & $F_{p,q}^\alpha(\mathbb{R}^n)$ / §I

Let $\varphi_0 \in \mathcal{S}(\mathbb{R}^n)$ be as in (1.1) and let $\tilde{\varphi}^{(0)} := \varphi_0$ and $\tilde{\varphi}^{(j)} := \varphi^{(j)}$ for any $j \in \mathbb{N}$, where $\varphi^{(j)}$ is as in (1.2). Then

$$\sum_{j=0}^{\infty} \tilde{\varphi}^{(j)}(x) = 1, \quad \forall x \in \mathbb{R}^n.$$

► The **inhomogeneous** Besov space $B_{p,q}^\alpha(\mathbb{R}^n)$ & Triebel-Lizorkin space $F_{p,q}^\alpha(\mathbb{R}^n)$ are defined via replacing $\mathcal{S}'_\infty(\mathbb{R}^n)$ and $\{\varphi^{(j)}\}_{j \in \mathbb{Z}}$ in $\dot{B}_{p,q}^\alpha(\mathbb{R}^n)$ & $\dot{F}_{p,q}^\alpha(\mathbb{R}^n)$, respectively, by $\mathcal{S}'(\mathbb{R}^n)$ (**Schwartz distributions**) and $\{\tilde{\varphi}^{(j)}\}_{j \in \mathbb{Z}_+}$. Moreover,

$$\|f\|_{F_{\infty,q}^\alpha(\mathbb{R}^n)} := \sup_{\substack{x \in \mathbb{R}^n \\ m \in \mathbb{Z}}} \left\{ 2^{mn} \int_{B(x, 2^{-m})} \sum_{j=\max\{m,0\}}^{\infty} |2^{j\alpha} (\tilde{\varphi}^{(j)} \hat{f})^\vee(y)|^q dy \right\}^{\frac{1}{q}}.$$

Hajłasz-Sobolev Spaces (1) / §I

- (\mathcal{X}, d, μ) : \mathcal{X} — nonempty set;
 d — quasi metric, namely, $d(x, y) \leq A[d(x, z) + d(z, y)]$;
 μ — regular Borel measure

- $p \in (1, \infty)$, $s \in (0, 1]$

- The **homogeneous fractional Hajłasz-Sobolev space** $\dot{M}^{s,p}(\mathcal{X})$ is defined to be the set of all measurable functions $f \in L^p_{\text{loc}}(\mathcal{X})$ for which there exist a $0 \leq g \in L^p(\mathcal{X})$ and a set $E \subset \mathcal{X}$ of measure zero such that, for any $x, y \in \mathcal{X} \setminus E$,

$$(1.3) \quad |f(x) - f(y)| \leq [d(x, y)]^s [g(x) + g(y)].$$

- Denote by $\mathcal{D}(f)$ the class of all non-negative Borel

Hajłasz-Sobolev Spaces (2) / §I

measurable functions g satisfying (1.3). Moreover, define

$$\|f\|_{\dot{M}^{s,p}(\mathcal{X})} := \inf_{g \in \mathcal{D}(f)} \{\|g\|_{L^p(\mathcal{X})}\}.$$

Let $M^{s,p}(\mathcal{X}) := L^p(\mathcal{X}) \cap \dot{M}^{s,p}(\mathcal{X})$ and, for any $f \in M^{s,p}(\mathcal{X})$, let

$$\|f\|_{M^{s,p}(\mathcal{X})} := \|f\|_{L^p(\mathcal{X})} + \|f\|_{\dot{M}^{s,p}(\mathcal{X})}.$$

Remarks:

- ▶ $\dot{M}^{1,p}(\mathcal{X})$ & $M^{1,p}(\mathcal{X})$ were introduced by Hajłasz [H96].
- ▶ $\dot{M}^{s,p}(\mathcal{X})$ & $M^{s,p}(\mathcal{X})$ when $s \in (0, 1)$ were introduced by Hu [Hu03] for subsets (fractals) of \mathbb{R}^n and Yang [Y03] for metric measure spaces.

Hajłasz-Sobolev Spaces (3) / §I

► It was proved in [H96] that

$$\dot{M}^{1,p}(\mathbb{R}^n) = \dot{W}^{1,p}(\mathbb{R}^n) = \dot{F}_{p,2}^1(\mathbb{R}^n)$$

and in [Y03] that, when $s \in (0, 1)$,

$$\dot{M}^{s,p}(\mathbb{R}^n) = \dot{F}_{p,\infty}^s(\mathbb{R}^n) \not\supseteq \dot{F}_{p,2}^s(\mathbb{R}^n).$$

(There exists a **gap** for **Triebel-Lizorkin spaces**.)

- [H96] **P. Hajłasz**, Sobolev spaces on an arbitrary metric space, **Potential Anal. 5 (1996), 403-415**.
- [Hu03] **J. Hu**, A note on Hajłasz-Sobolev spaces on fractals, **J. Math. Anal. Appl. 280 (2003), 91-101**.
- [Y03] **D. Yang**, New characterizations of Hajłasz-Sobolev spaces on metric spaces, **Sci. China Ser. A 46 (2003), 675-689**.

Fractional s –Hajłasz Gradient / §I

- [KYZ11] **P. Koskela, D. Yang & Y. Zhou**, Pointwise characterizations of Besov and Triebel-Lizorkin spaces and quasiconformal mappings, **Adv. Math. 226 (2011), 3579-3621**.

► **Definition.** Let $s \in (0, \infty)$ and u be a measurable function on \mathcal{X} . A sequence of nonnegative measurable functions, $\vec{g} := \{g_k\}_{k \in \mathbb{Z}}$, is called a **fractional s -Hajłasz gradient** of u if there exists $E \subset \mathcal{X}$ with $\mu(E) = 0$ such that, for any $k \in \mathbb{Z}$ and $x, y \in \mathcal{X} \setminus E$ satisfying $2^{-k-1} \leq d(x, y) < 2^{-k}$,

$$|u(x) - u(y)| \leq [d(x, y)]^s [g_k(x) + g_k(y)].$$

Denote by $\mathbb{D}^s(u)$ the collection of all fractional s -Hajłasz gradients of u .

$\dot{M}_{p,q}^s(\mathcal{X})$ & $\dot{N}_{p,q}^s(\mathcal{X})$ / §I

- The **homogeneous Hajlasz-Triebel-Lizorkin space** $\dot{M}_{p,q}^s(\mathcal{X})$ is defined to be the space of all measurable functions u such that

$$\|u\|_{\dot{M}_{p,q}^s(\mathcal{X})} := \inf_{\vec{g} \in \mathbb{D}^s(u)} \left\| \left\{ g_j \right\}_{j \in \mathbb{Z}} \right\|_{\ell^q} \Big\|_{L^p(\mathcal{X})} < \infty.$$

- The **homogeneous Hajlasz-Besov space** $\dot{N}_{p,q}^s(\mathcal{X})$ is defined to be the space of all measurable functions u such that

$$\|u\|_{\dot{N}_{p,q}^s(\mathcal{X})} := \inf_{\vec{g} \in \mathbb{D}^s(u)} \left\| \left\{ \|g_j\|_{L^p(\mathcal{X})} \right\}_{j \in \mathbb{Z}} \right\|_{\ell^q} < \infty.$$

RD-Spaces (1) / §I

• A triple (\mathcal{X}, d, μ) : \mathcal{X} is a non-empty set, d a quasi-metric (usually, for simplicity, metric), and μ a regular Borel measure.

▶ A space of homogenous type of Coifman-Weiss: if μ -**doubling** ($\mu(B(x, 2r)) \lesssim \mu(B(x, r))$).

▶ An RD-space if μ is both doubling and **reverse-doubling** ($\mu(B(x, 2r)) \geq C_0 \mu(B(x, r))$ and $C_0 > 1$).

▶ There exist many examples of RD-spaces. Especially, all **connected** spaces of homogeneous type are RD-spaces.

RD-Spaces (2) / §I

- [HMY06] **Y. Han, D. Müller & D. Yang**, Littlewood-Paley characterizations for Hardy spaces on spaces of homogeneous type, **Math. Nachr.** **279 (2006)**, 1505-1537.
- [HMY08] **Y. Han, D. Müller & D. Yang**, A theory of Besov and Triebel-Lizorkin spaces on metric measure spaces modeled on Carnot-Carathéodory spaces, **Abstr. Appl. Anal.** **2008 Art. ID 893409**, 250 pp.
- [MY09] **D. Müller & D. Yang**, A difference characterization of Besov and Triebel-Lizorkin spaces on RD-spaces, **Forum Math.** **21 (2009)**, 259-298.
- **D. Yang & Y. Zhou**, New properties of Besov and Triebel-Lizorkin spaces on RD-spaces, **Manuscripta Math.** **134 (2011)**, 59-90.

Theorem 1.5 / §I

► [KYZ11] **Theorem**. Let \mathcal{X} be an RD-space with the upper dimension n .

(i) If $s \in (0, 1)$, $p \in (n/(n + s), \infty)$ and $q \in (n/(n + s), \infty]$, then $\dot{M}_{p,q}^s(\mathcal{X}) = \dot{F}_{p,q}^s(\mathcal{X})$.

(ii) If $s \in (0, 1)$, $p \in (n/(n + s), \infty)$ and $q \in (0, \infty]$, then $\dot{N}_{p,q}^s(\mathcal{X}) = \dot{B}_{p,q}^s(\mathcal{X})$.

- $\dot{F}_{p,q}^s(\mathcal{X})$ & $\dot{B}_{p,q}^s(\mathcal{X})$ were studied in [HMY08] & [MY09].

- Applications to the **invariance** under **quasiconformal mappings** of the certain function spaces.

► More recent related papers: **H. Koch, P. Koskela, E. Saksman & T. Soto** [JFA, 2014], **M. Bonk, E. Saksman & T. Soto** [arXiv: 1411.5906 or **Indiana Univ. Math. J.** (to appear) or **D. Yang, W. Yuan & Y. Zhou** [JGA, 2017].

More Papers on $\dot{M}_{p,q}^s(\mathcal{X})$ & $\dot{N}_{p,q}^s(\mathcal{X})$ / §I

- ▶ **A. Gogatishvili, P. Koskela & Y. Zhou**, Characterizations of Besov and Triebel-Lizorkin spaces on metric measure spaces, **Forum Math. 25 (2013), 787-819.**
- ▶ **T. Heikkinen & H. Tuominen**, Approximation by Hölder functions in Besov and Triebel-Lizorkin spaces, **Constr. Approx. 44 (2016), 455-482.**
- ▶ **T. Heikkinen, L. Ihnatsyeva & H. Tuominen**, Measure density and extension of Besov and Triebel-Lizorkin functions, **J. Fourier Anal Appl. 22 (2016), 334-382.**
- ▶ **T. Heikkinen, P. Koskela & H. Tuominen**, Approximation and quasicontinuity of Besov and Triebel-Lizorkin functions, **Trans. Amer. Math. Soc. 369 (2017), 3547-3573.**

§II. **Ball average** characterizations of
second order Sobolev spaces

Theorem of [AMV12] / §II

► For any $t \in (0, \infty)$, $g \in L^1_{\text{loc}}(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$, let

$$B_t(g)(x) := \frac{1}{|B(x,t)|} \int_{B(x,t)} g(y) dy.$$

► ([AMV12]) Let $p \in (1, \infty)$. Then $f \in W^{2,p}(\mathbb{R}^n)$ if and only if $f \in L^p(\mathbb{R}^n)$ and there exists $g \in L^p(\mathbb{R}^n)$ such that

$$\mathcal{G}(f, g)(\cdot) := \left\{ \int_0^\infty \left| \frac{B_t(f)(\cdot) - f(\cdot)}{t^2} - B_t(g)(\cdot) \right|^2 \frac{dt}{t} \right\}^{\frac{1}{2}} \in L^p(\mathbb{R}^n).$$

► theory of **Vector-valued C-Z operators**, fine estimates

► [AMV12] **R. Alabern, J. Mateu & J. Verdera**, A new characterization of Sobolev spaces on \mathbb{R}^n , **Math. Ann. 354 (2012), 589-626.**

Lusin-Area Funct. Charact. (1) / §II

► ([HYY15]) (i) If $p \in [2, \infty)$, then $f \in W^{2,p}(\mathbb{R}^n)$ if and only if $f \in L^p(\mathbb{R}^n)$ and there exists $g \in L^p(\mathbb{R}^n)$ such that

$$\mathcal{S}(f, g)(\cdot) := \left\{ \int_0^\infty \int_{B(\cdot, t)} \left| \frac{B_t(f)(y) - f(y)}{t^2} - B_t(g)(y) \right|^2 dy \frac{dt}{t^{n+1}} \right\}^{\frac{1}{2}} \in L^p(\mathbb{R}^n).$$

(ii) If $p \in (1, 2)$ and $n \in \{1, 2, 3\}$, then $f \in W^{2,p}(\mathbb{R}^n)$ if and only if $f \in L^p(\mathbb{R}^n)$ and there exists $g \in L^p(\mathbb{R}^n)$ such that $\mathcal{S}(f, g) \in L^p(\mathbb{R}^n)$.

- [HYY15] **Z. He, D. Yang & W. Yuan**, Littlewood-Paley characterizations of second-order Sobolev spaces via averages on balls, **Canadian Math. Bull.** **59 (2016), 104-108.**

Lusin-Area Funct. Charact. (2) / §II

▶ ([DLYY]) (i) Let $n \in [4, \infty) \cap \mathbb{N}$ and $p \in (\frac{2n}{4+n}, 2)$. Then $f \in W^{2,p}(\mathbb{R}^n)$ if and only if $f \in L^p(\mathbb{R}^n)$ and there exists $g \in L^p(\mathbb{R}^n)$ such that $\mathcal{S}(f, g) \in L^p(\mathbb{R}^n)$.

(ii) Let $n \in [5, \infty) \cap \mathbb{N}$ and $p \in (1, \frac{2n}{4+n})$. Then the conclusion of (i) does not hold true.

▶ In (i), if $n = 4$, then $p \in (1, 2)$. The conclusion of (i) is **near sharp**.

▶ Instead of **a** vector-valued C-Z operator, use **a series of** vector-valued C-Z operators

• [DLYY] **F. Dai, J. Liu, D. Yang & W. Yuan**, Littlewood-Paley characterizations of fractional Sobolev spaces via averages on balls, **Proc. Roy. Soc. Edinburgh Sect. A.** (to appear).

\mathcal{G}_λ^* Characterization (1) / §II

► ([HYY15]) (i) If $p \in [2, \infty)$ and $\lambda \in (1, \infty)$, then
 $f \in W^{2,p}(\mathbb{R}^n) \iff f \in L^p(\mathbb{R}^n)$ and $\exists g \in L^p(\mathbb{R}^n)$ such that

$$\mathcal{G}_\lambda^*(f, g)(\cdot) := \left\{ \int_0^\infty \int_{\mathbb{R}^n} \left| \frac{B_t(f)(y) - f(y)}{t^2} - B_t(g)(y) \right|^2 \right. \\ \left. \times \left(\frac{t}{t + |\cdot - y|} \right)^{\lambda n} dy \frac{dt}{t^{n+1}} \right\}^{\frac{1}{2}} \in L^p(\mathbb{R}^n).$$

(ii) If $p \in (1, 2)$, $\lambda \in (2/p, \infty)$ and $n \in \{1, 2, 3\}$, then
 $f \in W^{2,p}(\mathbb{R}^n) \iff f \in L^p(\mathbb{R}^n)$ and $\exists g \in L^p(\mathbb{R}^n)$ such that
 $\mathcal{G}_\lambda^*(f, g) \in L^p(\mathbb{R}^n)$.

\mathcal{G}_λ^* Characterization (2) / §II

▶ ([DLYY]) (i) Let $n \in [4, \infty) \cap \mathbb{N}$, $p \in (\frac{2n}{4+n}, 2)$ and $\lambda \in (2/p, \infty)$. Then $f \in W^{2,p}(\mathbb{R}^n)$ if and only if $f \in L^p(\mathbb{R}^n)$ and there exists $g \in L^p(\mathbb{R}^n)$ such that $\mathcal{G}_\lambda^*(f, g) \in L^p(\mathbb{R}^n)$.

(ii) Let $n \in [5, \infty) \cap \mathbb{N}$, $p \in (1, \frac{2n}{4+n})$ and $\lambda \in (2/p, \infty)$. Then the conclusion of (i) does not hold true.

▶ In (i), if $n = 4$, then $p \in (1, 2)$. The conclusion of (i) is **near sharp**.

▶ It is still unclear on the endpoint case $p = \frac{2n}{4+n}$.

Pointwise Characterization (1) / §II

► ([DGY15]) Let $p \in (1, \infty)$. Then $f \in W^{2,p}(\mathbb{R}^n)$

$\iff f \in L^p(\mathbb{R}^n)$ and $\exists 0 \leq g \in L^p(\mathbb{R}^n)$ and $C_0 > 0$ such that, for any $t \in (0, \infty)$ and almost every $x \in \mathbb{R}^n$,

$$|f(x) - B_t(f)(x)| \leq C_0 t^2 g(x)$$

$\iff f \in L^p(\mathbb{R}^n)$ and

$$\sup_{t \in (0, \infty)} \frac{\|f - B_t(f)\|_{L^p(\mathbb{R}^n)}}{t^2} =: C_1 < \infty.$$

• **Not known** for spaces of homogenous type.

• [DGY15] **F. Dai, A. Gogatishvili, D. Yang & W. Yuan,**

Characterizations of Sobolev spaces via averages on balls,

Nonlinear Anal. 128 (2015), 86-99.

Pointwise Characterization (2) / §II

► ([DGY15]) Let $p \in (1, \infty)$. Then $f \in W^{2,p}(\mathbb{R}^n)$

$\iff f \in L^p(\mathbb{R}^n)$ and $\exists 0 \leq g \in L^p(\mathbb{R}^n)$ and $C, \tilde{C} > 0$ such that, for any $t \in (0, \infty)$ and almost every $x \in \mathbb{R}^n$,

$$|B_t(f - B_{\tilde{C}t}(f))(x)| \leq Ct^2 g(x)$$

$\iff f \in L^p(\mathbb{R}^n)$ and $\exists 0 \leq g \in L^p(\mathbb{R}^n)$ and $c, C, \tilde{C} > 0$ such that, for any $t \in (0, \infty)$ and almost every $x \in \mathbb{R}^n$,

$$B_t(|f - B_{\tilde{C}t}(f)|)(x) \leq Ct^2 B_{ct}(g)(x).$$

► The second equivalence also holds true on **spaces of homogeneous type**.

Pointwise Characterization (3) / §II

► ([DGYY15]) Let $p \in (1, \infty)$, $q \in [1, p)$, $c \in (0, \infty)$ and $K \in (0, \infty]$. Then $f \in W^{2,p}(\mathbb{R}^n)$

$\iff f \in L^p(\mathbb{R}^n)$ and

$$f_{c,q}^{\#,K}(\cdot) := \sup_{t \in (0,K)} t^{-2} \{B_t(|f - B_{ct}(f)|^q)(\cdot)\}^{1/q} \in L^p(\mathbb{R}^n).$$

A Key Lemma / §II

Let $\varphi \in \mathcal{S}(\mathbb{R}^n)$ and $\tilde{C} \in (0, \infty)$ be a constant. Then

$$\lim_{t \rightarrow 0^+} \frac{\varphi - B_t(\varphi)}{t^2} = -\frac{1}{2(n+2)} \Delta \varphi$$

and

$$\lim_{t \rightarrow 0^+} B_t \left(\frac{\varphi - B_{\tilde{C}t}(\varphi)}{t^2} \right) (\cdot) = -\frac{\tilde{C}^2}{2(n+2)} \Delta \varphi(\cdot)$$

with convergence in $\mathcal{S}(\mathbb{R}^n)$.

Further Results / §II

► For any $\ell \in \mathbb{N}$, $t \in (0, \infty)$, $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$, let

$$B_{\ell,t}(f)(x) := -\frac{2}{\binom{2\ell}{\ell}} \sum_{j=1}^{\ell} (-1)^j \binom{2\ell}{\ell-j} B_{jt}(f)(x).$$

(Binomial coefficients; Observe that $B_{1,t}(f) = B_t(f)$.)

► All aforementioned characterizations of $W^{2,p}(\mathbb{R}^n)$ via pointwise inequalities remain true for $W^{2\ell,p}(\mathbb{R}^n)$, with $\ell \in \mathbb{N}$ and $p \in (1, \infty)$, if we replace $B_t(f)$ by $B_{\ell,t}(f)$ therein.

► ([CYYZ]) Also true for Morrey-Sobolev spaces.

• [CYYZ] **D.-C. Chang, D. Yang, W. Yuan & J. Zhang**, Some recent developments of high order Sobolev-type spaces, **J. Nonlinear Convex Anal.** **17** (2016), 1831-1865.

Open Questions / §II

- ▶ On spaces of homogeneous type (or even **smooth domains** of \mathbb{R}^n), whether or not these Sobolev spaces **coincide**? (We now have **several different** definitions.)
- ▶ On spaces of homogeneous type, whether or not fractional Sobolev spaces **contain** the known Hajlasz-Sobolev spaces or the known Newton-Sobolev spaces?
- ▶ For analysis on metric measure spaces, any **applications**?



§III. **Ball average** characterizations of
second order Besov and
Triebel-Lizorkin spaces

Littlewood-Paley Characterization (1) / §III

► ([YYZ13, DGYY15]) Let $\alpha \in (0, 2)$ and $q \in (1, \infty]$.

(i) If $p \in (1, \infty)$, then $f \in F_{p,q}^\alpha(\mathbb{R}^n)$ if and only if $f \in L^p(\mathbb{R}^n)$ and

$$\left\| \left\{ 2^{k\alpha} |f - B_{2^{-k}}(f)| \right\}_{k \in \mathbb{Z}_+} \right\|_{\ell^q} \Big\|_{L^p(\mathbb{R}^n)} < \infty.$$

(ii) If $p = \infty$, then $f \in F_{\infty,q}^\alpha(\mathbb{R}^n)$ if and only if $f \in C(\mathbb{R}^n)$ and

$$\sup_{\substack{\ell \in \mathbb{Z} \\ x \in \mathbb{R}^n}} \left[B_{2^{-\ell}} \left(\sum_{k \geq \max\{\ell, 0\}} 2^{k\alpha q} |f - B_{2^{-k}}(f)|^q \right) (x) \right]^{\frac{1}{q}} < \infty.$$

► $C(\mathbb{R}^n)$: the space of all **uniformly continuous bounded** functions

Littlewood-Paley Characterization (2) / §III

► ([YYZ13, DGYY15]) Let $\alpha \in (0, 2)$, $p \in (1, \infty]$ and $q \in (0, \infty]$. Then $f \in B_{p,q}^\alpha(\mathbb{R}^n)$ if and only if $f \in L^p(\mathbb{R}^n)$ when $p < \infty$, or $f \in C(\mathbb{R}^n)$ when $p = \infty$, and

$$\left\{ \sum_{j=0}^{\infty} 2^{j\alpha q} \|f - B_{2^{-j}}(f)\|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}} < \infty.$$

- [YYZ13] **D. Yang, W. Yuan & Y. Zhou**, A new characterization of Triebel-Lizorkin spaces on \mathbb{R}^n , **Publ. Mat. 57 (2013), 57-82**.
- [DGYY15] **F. Dai, A. Gogatishvili, D. Yang & W. Yuan**, Characterizations of Besov and Triebel-Lizorkin spaces via averages on balls, **J. Math. Anal. Appl. 433 (2016), 1350-1368**.

Littlewood-Paley Characterization (3) / §III

▶ Lusin-area type function: For any $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$,

$$\mathcal{A}_r(f)(x) := \left\{ \sum_{k=1}^{\infty} 2^{k\alpha q} [B_{2^{-k}}(|f - B_{2^{-k}}(f)|^r)(x)]^{\frac{q}{r}} \right\}^{\frac{1}{q}}.$$

▶ ([CLYY15]) Let $\alpha \in (0, 2)$, $p \in (1, \infty)$, $q \in (1, \infty]$ and $r \in [1, q)$. Then $f \in F_{p,q}^\alpha(\mathbb{R}^n)$ if and only if $f \in L^p(\mathbb{R}^n)$ and $\mathcal{A}_r(f) \in L^p(\mathbb{R}^n)$.

▶ [CLYY15] **D.-C. Chang, J. Liu, D. Yang & W. Yuan**, Littlewood-Paley characterizations of Hajlasz-Sobolev and Triebel-Lizorkin spaces via averages on balls, **Potential Anal.** **46 (2017), 22-259.**

Littlewood-Paley Characterization (4) / §III

- ▶ ([CLYY15]) Let $\alpha \in (0, 2)$ and $p \in (1, \infty)$, $q \in (1, \infty]$.
 - (i) If $f \in L^p(\mathbb{R}^n)$ and $\mathcal{A}_q(f) \in L^p(\mathbb{R}^n)$, then $f \in F_{p,q}^\alpha(\mathbb{R}^n)$.
 - (ii) If $p \in [q, \infty)$ and $\alpha \in (0, 2)$, or $p \in (1, q)$ and $\alpha \in (n(1/p - 1/q), 2)$, then $f \in F_{p,q}^\alpha(\mathbb{R}^n)$ implies that $f \in L^p(\mathbb{R}^n)$ and $\mathcal{A}_q(f) \in L^p(\mathbb{R}^n)$.

- ▶ In case when $q = 2$ and $\alpha < 1$, i. e., $F_{p,2}^\alpha(\mathbb{R}^n) = W^{\alpha,p}(\mathbb{R}^n)$, then (ii) is not true when $\alpha < n(1/p - 1/2)$. But, what happens when $\alpha = n(1/p - 1/2)$?

Littlewood-Paley Characterization (5) / §III

- ▶ Part of aforementioned results are also true for $B_{p,q}^{\alpha,\tau}(\mathbb{R}^n)$ & $F_{p,q}^{\alpha,\tau}(\mathbb{R}^n)$ (see [ZSY]) and Triebel-Lizorkin-Morrey spaces $\mathcal{E}_{u,q,p}^{\alpha}(\mathbb{R}^n)$ (see [ZZYH]).
- ▶ [ZSY] **C. Zhuo, W. Sickel, D. Yang & W. Yuan**, Characterizations of Besov-type and Triebel-Lizorkin-type spaces via averages on balls, **Canad. Math. Bull.** 60 (2017), 655-672.
- ▶ [ZZYH] **J. Zhang, C. Zhuo, D. Yang and Z. He**, Littlewood-Paley characterizations of Triebel-Lizorkin-Morrey spaces via ball averages, **Nonlinear Anal.** 150 (2017), 76-103.
- ▶ A new idea is to introduce the **local** Hardy-Littlewood maximal function.

Pointwise Characterization (1) / §III

► ([YY15]) Let $\alpha \in (0, \infty)$ and $f \in L^1_{\text{loc}}(\mathbb{R}^n)$. A sequence $\vec{g} := \{g_j\}_{j \geq 0}$ of non-negative measurable functions is called an α -order Hajłasz type gradient sequence of f if, for each j , there exists a set $E_j \subset \mathbb{R}^n$ with measure zero such that

$$|f(x) - B_{2^{-j}} f(x)| \leq 2^{-j\alpha} g_j(x), \quad \forall x \in \mathbb{R}^n \setminus E_j.$$

Each g_j satisfying the above is called an α -order Hajłasz type gradient of f at level j .

► [YY15] **D. Yang & W. Yuan**, Pointwise characterizations of Besov and Triebel-Lizorkin spaces in terms of averages on balls, **Trans. Amer. Math. Soc.** **369** (2017), 7631-7655.

Pointwise Characterization (2) / §III

► ([YY15]) Let $\alpha \in (0, 2)$ and $p, q \in (1, \infty]$. Then $f \in F_{p,q}^\alpha(\mathbb{R}^n)$ if and only if $f \in L^p(\mathbb{R}^n)$ when $p \in (0, \infty)$ or $f \in C(\mathbb{R}^n)$ when $p = \infty$, and there exists an **α -order Hajlasz type gradient sequence** $\vec{g} := \{g_k\}_{k=0}^\infty$ of f such that

$$\left\| \left\{ \sum_{k=0}^{\infty} 2^{k\alpha q} |g_k|^q \right\}^{1/q} \right\|_{L^p(\mathbb{R}^n)} < \infty, \quad p < \infty$$

and

$$\sup_{\substack{\ell \in \mathbb{Z} \\ x \in \mathbb{R}^n}} \left[B_{2^{-\ell}} \left(\sum_{k \geq \max\{\ell, 0\}} 2^{k\alpha q} |g_k|^q \right) (x) \right]^{1/q} < \infty, \quad p = \infty.$$

Pointwise Characterization (3) / §III

► ([YY15]) Let $\alpha \in (0, 2)$, $p \in (1, \infty]$ and $q \in (0, \infty]$. Then $f \in B_{p,q}^\alpha(\mathbb{R}^n)$ if and only if $f \in L^p(\mathbb{R}^n)$ when $p \in (0, \infty)$, or $f \in C(\mathbb{R}^n)$ when $p = \infty$, and there exists an **α -order Hajlasz type gradient sequence** $\vec{g} := \{g_k\}_{k=0}^\infty$ of f such that

$$\left\{ \sum_{k=0}^{\infty} 2^{k\alpha q} \|g_k\|_{L^p(\mathbb{R}^n)} \right\}^{1/q} < \infty.$$

► Let $\ell \in \mathbb{N}$. All aforementioned pointwise characterizations of $B_{p,q}^\alpha(\mathbb{R}^n)$ and $F_{p,q}^\alpha(\mathbb{R}^n)$ remain true when $\alpha \in (0, 2\ell)$ if we replace $\{f - B_{2^{-j}}(f)\}_j$ by $\{f - B_{\ell, 2^{-j}}(f)\}_j$.

► Applications?



§IV. Further remarks

Newton Spaces (1) / §IV

▶ **N. Shanmugalingam**, Newtonian spaces: An extension of Sobolev spaces to metric measure spaces, **Rev. Mat. Iberoamericana** **16** (2000), 243-279.

▶ Defined via **upper gradients** (J. Heinonen & P. Koskela [Acta Math., 1998]; P. Koskela & P. MacManus [Studia Math., 1998]). Instead of straight lines by **curves** γ :

$$|f(\gamma(a)) - f(\gamma(b))| \leq \int_{\gamma} g ds.$$

▶ Advantage: **Strong locality** (If a function is **constant on a measurable set**, then we can take **upper gradient to be zero almost everywhere on that set**; however, we cannot take the **Hajlasz gradient to be zero almost everywhere on that set**.)

▶ **N. Shanmugalingam, D. Yang & W. Yuan**, Newton-Besov spaces and Newton-Triebel-Lizorkin spaces, **Positivity** **19** (2015), 177-220.

Newton Spaces (2) / §IV

- ▶ **Newtonian-type Orlicz-Sobolev** spaces on metric measure spaces
 - **H. Tuominen**, Orlicz-Sobolev spaces on metric measure spaces, **Dissertation, University of Jyväskylä, Jyväskylä, 2004. Ann. Acad. Sci. Fenn. Math. Diss. No. 135 (2004), 86 pp.**
- ▶ **Hajlasz-type Orlicz-Sobolev** spaces and **Newtonian-type Orlicz-Sobolev** spaces
 - **T. Ohno & T. Shimomura**, Musielak-Orlicz-Sobolev spaces on metric measure spaces, **Czechoslovak Math. J. 65 (140) (2015), 435-474.**

Sphere Average Charact. / §IV

- ▶ **P. Hajłasz & Z. Liu**, A Marcinkiewicz integral type characterization of the Sobolev space, **Publ. Mat. 61 (2017), 83-104.**

- Let $p \in (1, \infty)$. Then $f \in W^{1,p}(\mathbb{R}^n)$ if and only if $f \in L^p(\mathbb{R}^n)$ and

$$\left[\int_0^\infty \left| f(\cdot) - \frac{1}{|S(\cdot, t)|} \int_{S(\cdot, t)} f(y) d\sigma(y) \right|^2 \frac{dt}{t^3} \right]^{1/2} \in L^p(\mathbb{R}^n),$$

where $S(x, t)$ denotes the sphere centered at x with the radius t . (**Not so useful for metric measure spaces**)

Weighted Sobolev Spaces / §IV

- A new and simplified proof of the characterization of $W^{1,p}(\mathbb{R}^n)$ with $p \in (1, \infty)$ ([Theorem 1, AMV12])
- ▶ **S. Sato**, Littlewood-Paley operators and Sobolev spaces, **Illinois J. Math.** **58 (2014), 1025-1039.**
 - Generalize [Theorem 1, AMV12] to the weighted case: $W_w^{\alpha,p}(\mathbb{R}^n)$, $\alpha \in (0, 2)$, $p \in (1, \infty)$ and $w \in A_p(\mathbb{R}^n)$
- ▶ **S. Sato**, Littlewood-Paley equivalence and homogeneous Fourier multipliers, **Integral Equations Operator Theory** **87 (2017), 15-44.**
- ▶ **S. Sato**, Spherical square functions of Marcinkiewicz type with Riesz potentials, **Arch. Math. (Basel)** **108 (2017), 415-426.**

Generalized Means / §IV

- ▶ Let Φ be a **bounded radial** function on \mathbb{R}^n with **compact support** satisfying $\int_{\mathbb{R}^n} \Phi(x) dx = 1$.
- ▶ Generalized means:

$$G_t(g)(x) := \int_{\mathbb{R}^n} \frac{1}{t^n} \Phi\left(\frac{x-y}{t}\right) g(y) dy.$$

If let $\Phi := \frac{1}{|B(\vec{0}_n, 1)|} \chi_{B(\vec{0}_n, 1)}$, then $G_t(g) = B_t(g)$.

- **S. Sato, F. Wang, D. Yang & W. Yuan**, Generalized Littlewood-Paley characterizations of fractional Sobolev spaces, **Commun. Contemp. Math.** (to appear). [only $\alpha \in (0, 2]$]
- **Y. Zhang, D.-C. Chang & D. Yang**, Generalized Littlewood-Paley characterizations of Triebel-Lizorkin spaces, **J. Nonlinear Convex Anal.** 18 (2017), 1171-1190. [only $\alpha \in (0, 2)$]

Morrey-Sobolev Spaces / §IV

- **Morrey**-Sobolev Spaces on Metric Measure Spaces

- ▶ Let $0 < p \leq q \leq \infty$. Recall that the **Morrey space** $\mathcal{M}_p^q(\mathcal{X})$ is defined to be the space of all measurable functions f on \mathcal{X} such that

$$\|f\|_{\mathcal{M}_p^q(\mathcal{X})} := \sup_{B \subset \mathcal{X}} [\mu(B)]^{1/q-1/p} \left[\int_B |f(x)|^p d\mu(x) \right]^{1/p} < \infty,$$

where the supremum is taken over all balls B in \mathcal{X} .

- ▶ Replace $L^p(\mathcal{X})$ by $\mathcal{M}_p^q(\mathcal{X})$
- ▶ **Y. Lu, D. Yang & W. Yuan**, Morrey-Sobolev spaces on metric measure spaces, **Potential Anal. 41 (2014), 215-243.**


Haroske and Triebel / §IV

- Triebel [T10] introduced the **higher** version of Hajlasz-Sobolev spaces on \mathbb{R}^n via **higher differences**, and some very interesting applications are given in [T11] and [HT11]:
 - ▶ [HT11] **D. D. Haroske & H. Triebel**, Embeddings of function spaces: a criterion in terms of differences, **Complex Var. Elliptic Equ.** **56 (2011), 931-944.**
 - ▶ [T10] **H. Triebel**, Sobolev-Besov spaces of measurable functions, **Studia Math.** **201 (2010), 69-86.**
 - ▶ [T11] **H. Triebel**, Limits of Besov norms, **Arch. Math.** **96 (2011), 169-175.**

Sobolev Spaces Associated with Operators / §IV

- ▶ **L. Yan & D. Yang**, New Sobolev spaces via generalized Poincaré inequalities on metric measure spaces, **Math. Z.** **255** (2007), 133-159.
- ▶ **S. Hofmann, S. Mayboroda & A. McIntosh**, Second order elliptic operators with complex bounded measurable coefficients in L^p , Sobolev and Hardy spaces, **Ann. Sci. École Norm. Sup. (4)** **44** (2011), 723-800.
- ▶ **F. Bernicot, T. Coulhon & F. Dorothee**, Sobolev algebras through heat kernel estimates, **J. Éc. polytech. Math.** **3** (2016), 99-161.
- ▶ **J. Zhang, D.-C. Chang & D. Yang**, Characterizations of Sobolev spaces associated to operators satisfying off-diagonal estimates on balls, **Math. Methods Appl. Sci.** **40** (2017), 2907-2929.

Lecture Notes in Math. 2182, 2017 / §IV



Thank you for your attention.